

Efficient Down-Regulation of Cyclin A-Associated Activity and Expression in Suspended Primary Keratinocytes Requires p21^{Cip1}

Paul Hauser,¹ Le Ma,¹ Deepak Agrawal,¹ Eric Haura,² W. Douglas Cress,¹ and W. Jackson Pledger¹

¹Molecular Oncology Program and ²Experimental Therapeutics Program, H. Lee Moffitt Cancer Center and Research Institute, Department of Interdisciplinary Oncology, University of South Florida School of Medicine, Tampa FL

Abstract

When suspended in methylcellulose, primary mouse keratinocytes cease proliferation and differentiate. Suspension also reduces the activity of the cyclin-dependent kinase cdk2, an important cell cycle regulatory enzyme. To determine how suspension modulates these events, we examined its effects on wild-type keratinocytes and keratinocytes nullizygous for the cdk2 inhibitor p21^{Cip1}. After suspension of cycling cells, amounts of cyclin A (a cdk2 partner), cyclin A mRNA, and cyclin A-associated activity decreased much more rapidly in the presence than in the absence of p21^{Cip1}. Neither suspension nor p21^{Cip1} status affected the stability of cyclin A mRNA. Loss of p21^{Cip1} reduced the capacity of suspended cells to growth arrest, differentiate, and accumulate p27^{Kip1} (a second cdk2 inhibitor) and affected the composition of E2F DNA binding complexes. Cyclin A-cdk2 complexes in suspended p21^{+/+} cells contained p21^{Cip1} or p27^{Kip1}, whereas most of the cyclin A-cdk2 complexes in p21^{-/-} cells lacked p27^{Kip1}. Ectopic expression of p21^{Cip1} allowed p21^{-/-} keratinocytes to efficiently down-regulate cyclin A and differentiate when placed in suspension. These findings show that p21^{Cip1} mediates the effects of suspension on numerous processes in primary keratinocytes including cdk2 activity, cyclin A expression, cell cycle progression, and differentiation.

Introduction

Most mammalian cell types grow in an adhesion-dependent manner both *in vivo* and *in vitro*. Oncogenic transformation overrides the need for attachment for cell proliferation, and anchorage-independent growth correlates strongly with tumorigenicity. When deprived of a substratum, nontransformed cells typically arrest in G₁, and studies addressing this phenomenon have focused on the cyclin-dependent kinases (CDKs) and their

regulatory subunits, the cyclins (1). Complexes containing the D cyclins and cdk4, cyclin E and cdk2, and cyclin A and cdk2 become active in mid-G₁, in late G₁, and at G₁-S, respectively (2). All complexes phosphorylate the retinoblastoma (Rb) protein, an event required for S phase entry (3). When hyperphosphorylated by cdk4 and cdk2, Rb no longer functions as a transcriptional repressor, and numerous gene products (including many involved in DNA replication) are expressed (4, 5). Events that regulate the activity of cdk4 and cdk2 include the periodic synthesis and destruction of the cyclins. In addition, proteins such as p21^{Cip1} and p27^{Kip1} interact with and inhibit the activity of complexes containing cdk4 or cdk2 and their cyclin partners (6, 7). The overall abundance of p21^{Cip1} and p27^{Kip1} varies during the cell cycle, and sequestration of p21^{Cip1} and p27^{Kip1} by D cyclin-cdk4 complexes facilitates cdk2 activation.

Most notably, nontransformed cells do not express cyclin A, and consequently do not form cyclin A-cdk2 complexes, when cultured in suspension (8–18). Conversely, cyclin A is readily detectable in cells that grow in suspension as a result of enforced expression of E7, c-Myc, activated Ras, cyclin D1, or integrin-linked kinase (11, 13, 15–17, 19, 20). When ectopically expressed, cyclin A allows or contributes to the anchorage-independent proliferation of NRK, Rat 1a, and ER-1-2 fibroblasts (8, 11, 13). Collectively, these studies identify cyclin A as a key target of suspension-induced growth inhibitory pathways.

Transcriptional repression of the cyclin A promoter accounts (at least in part) for the absence of cyclin A from suspended cells (10, 14, 21, 22). Repression occurs when Rb sequesters a CCAAT-binding protein required for transactivation and/or when Rb or the Rb-related protein p107 interacts with a variant E2F site in the cyclin A promoter. In both instances, Rb and p107 lose their capacity to repress *cyclin A* gene transcription when hyperphosphorylated by cdk4 and cdk2. Suspension inhibits cyclin E-cdk2 activity in most cell types and cdk4 activity in some cell types; as a result, Rb and p107 are typically hypophosphorylated in suspended cells (9–12, 15–17, 19, 23, 24). Suspension does not alter the abundance of cyclin E or cdk2 or the assembly of cyclin E-cdk2 complexes; however, more cyclin E-cdk2 complexes contain p21^{Cip1} or p27^{Kip1} in suspended cells than in attached cycling cells (9, 19, 23, 24). p21^{Cip1} and p27^{Kip1} often accumulate in suspended cells, and in some instances, the absence of the D cyclins precludes the sequestration of p21^{Cip1} and p27^{Kip1} by D cyclin-cdk4 complexes (9, 10, 12, 16, 19, 20, 23, 24). Thus, suspension increases the availability of p21^{Cip1} and p27^{Kip1}, which results

Received 9/15/03; revised 12/30/03; accepted 1/8/04.

Grant support: This work was supported by the Corner-Couch Endowed Chair for Cancer Research (W. J. P.) and NIH Grants CA73780 (W. J. P.) and CA90489 (W. D. C.).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked advertisement in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Requests for reprints: W.J. Pledger, H. Lee Moffitt Cancer Center and Research Institute, 12902 Magnolia Drive, Tampa FL 33612. Phone: (813) 979-3887; Fax: (813) 979-3393. E-mail: pledgerw@moffitt.usf.edu

Copyright © 2004 American Association for Cancer Research.

in cyclin E-cdk2 inactivation, Rb hypophosphorylation, and ultimately, the repression of *cyclin A* gene transcription. Suspension may also inhibit cyclin A expression by processes other than those described above (9, 11).

Most previous studies examined the effects of suspension on cells stimulated from quiescence (8–16, 22, 23). In contrast, our studies (18) examined the effects of suspension on growing cells, which express active cyclin A-cdk2 complexes at the time of suspension. Our studies were performed on primary mouse keratinocytes, which cease proliferation and differentiate when placed in a semisolid medium (25). This system allows us to determine how suspension affects cyclin-CDK activity and how changes in cyclin-CDK activity affect both cell cycle progression and differentiation. We found that p27^{Kip1} accumulated in suspended keratinocytes and interacted with cyclin A-cdk2 complexes and that amounts of cyclin A and cyclin A-associated activity declined (18). Suspension of cells in the presence of p27^{Kip1} antisense oligonucleotides, and thus in the absence of p27^{Kip1} accumulation, did not prevent the loss of cyclin A or cyclin A-associated activity or cell cycle arrest; however, the differentiation-specific marker keratin 1 (K1) was not induced.

On the basis of the results of our previous report (18), we suggest that p27^{Kip1} promotes keratinocyte differentiation by actions unrelated to the inhibition of cdk2 activity. We also suggest that p27^{Kip1} (or p27^{Kip1} accumulation) contributes to but is not required for cdk2 inactivation and growth arrest in suspended cells. To identify additional mediators of these events, we examined the effects of suspension on keratinocytes lacking p21^{Cip1}. We show that efficient down-regulation of cyclin A and cyclin A-associated activity in suspended keratinocytes requires p21^{Cip1} as do growth arrest, differentiation, and maximal accumulation of p27^{Kip1}. Determination of how suspension abrogates cell cycle controls in normal keratinocytes may be relevant to studies of cancer cells, which typically grow in suspension.

Results

Requirement for p21^{Cip1} for Efficient Down-Regulation of Cyclin A, Cyclin A Messenger RNA, and Cyclin A-Associated Activity in Suspended Keratinocytes

To assess the need for p21^{Cip1} for suspension-induced decreases in cyclin A expression and cyclin A-associated activity, primary keratinocytes derived from wild-type and p21^{Cip1}-null mice were suspended in methylcellulose and harvested at various times thereafter. At the time of suspension, keratinocytes in both populations were actively cycling (data not shown). Amounts of cyclin A were determined by Western blotting of cell extracts, and kinase activity was assayed in cyclin A immune complexes using histone H1 as substrate. Although cyclin A associates with both cdk2 and the mitotic CDK cdc2 (26), p21^{Cip1} selectively inhibits the activity of cdk2-containing complexes (27). It is noted that p21^{Cip1} preferentially interacts with cyclin A-cdk2 complexes rather than cyclin A or cdk2 alone (27).

Cyclin A was expressed and cyclin A-associated kinases were active in attached p21^{+/+} keratinocytes and, to a greater extent, in attached p21^{+/+} keratinocytes (Fig. 1, A and B, lanes 1 and 6). Cyclin A-associated activity decreased within

3 h of suspension of p21^{+/+} keratinocytes and was barely detectable at 6 h and times thereafter (Fig. 1B, lanes 1–5), even when the blots were overexposed (lanes 11–15). Amounts of cyclin A declined more slowly; cyclin A abundance was modestly reduced 6 h after suspension and substantially reduced 12 h after suspension of p21^{+/+} cells (Fig. 1A, lanes 1–5). Importantly, cyclin A was detectable in cells suspended for 6–12 h, whereas cyclin A-associated activity was not. This indicates that factors other than (or in addition to) loss of protein account for loss of activity during this time period.

In contrast to p21^{+/+} keratinocytes, p21^{-/-} keratinocytes expressed comparable amounts of cyclin A or cyclin A-associated activity whether attached to plates or suspended for 12 h (Fig. 1A, lanes 6–10, and Fig. 1B, lanes 6–9). Amounts of both, however, eventually declined (Fig. 1B, lane 10, and data not shown). The data in Fig. 1, A and B, indicate that loss of p21^{Cip1} greatly delays the suspension-induced down-regulation of cyclin A and cyclin A-associated activity in primary keratinocytes.

The effects of suspension on cyclin A mRNA abundance in p21^{+/+} and p21^{-/-} keratinocytes were similar to its effects on cyclin A protein abundance. After suspension, amounts of cyclin A mRNA decreased rapidly and substantially in p21^{+/+} keratinocytes (Fig. 1C, lanes 1–5), but remained elevated in p21^{-/-} keratinocytes for 12 h (lanes 6–10). When p21^{+/+} and p21^{-/-} keratinocytes were suspended in the presence of the mRNA synthesis inhibitor DRB, the cyclin A transcript decayed with similar kinetics in both populations (lanes 11–18). Thus, differences in cyclin A mRNA stability do not account for the differences in cyclin A mRNA abundance in suspended p21^{+/+} versus p21^{-/-} keratinocytes. Amounts of cyclin A mRNA also decreased with similar kinetics in p21^{+/+} keratinocytes suspended in the presence (lanes 11–14) and absence (lanes 2–5) of DRB. By default, these data suggest that loss of p21^{Cip1} prevents suspension-induced decreases in the activity of the cyclin A promoter. This premise is consistent with previous findings showing transcriptional repression of the *cyclin A* gene in nonadherent fibroblasts (10, 14, 21, 22).

To ensure that p21^{Cip1} status accounts for the different expression patterns of cyclin A in wild-type versus p21^{Cip1}-null keratinocytes, we ectopically expressed p21^{Cip1} in p21^{-/-} keratinocytes and amounts of cyclin A were determined. In these experiments, p21^{-/-} keratinocytes were infected with adenovirus without insert or adenovirus encoding human p21^{Cip1} (Ad-p21^{Cip1}). Cells were suspended in methylcellulose 19 h after infection and harvested 19 h later. For comparative purposes, p21^{+/+} keratinocytes were also examined (Fig. 2, lanes 1 and 2). The expression of human p21^{Cip1} in p21^{-/-} keratinocytes was verified by Western blotting (lanes 4 and 7). After suspension, amounts of cyclin A declined but were readily detectable in uninfected p21^{-/-} keratinocytes and p21^{-/-} keratinocytes infected with virus alone (compare lanes 3, 5, and 6). In contrast, cyclin A was not detected in p21^{-/-} keratinocytes infected with Ad-p21^{Cip1} and suspended for 19 h (lane 7). These findings show that nonadherent p21^{-/-} keratinocytes gain the capacity to efficiently down-regulate cyclin A when supplied with p21^{Cip1}. It is unclear why Ad-p21^{Cip1} more completely reduces cyclin A abundance in suspended keratinocytes (lane 7) than in attached keratinocytes (lane 4).




FIGURE 1. Amounts of cyclin A protein, mRNA, and associated activity in p21^{+/+} and p21^{-/-} keratinocytes suspended in methylcellulose. **A** and **B.** Proliferating p21^{+/+} and p21^{-/-} keratinocytes were detached from tissue cultures plates and suspended in methylcellulose as described in “Materials and Methods.” Cells were harvested at the times indicated. **A.** Amounts of cyclin A were determined by Western blotting. **B.** Cell extracts were immunoprecipitated with antibody to cyclin A, and cyclin A immune complexes were assayed for kinase activity using histone H1 as substrate. Lanes 1–10, results of a single exposure of the autoradiogram. Lanes 11–15, darker exposure of lanes 1–5, respectively. The darker exposure allows normalization of the activities at time 0 for p21^{+/+} (lane 11) and p21^{-/-} keratinocytes (lane 6). **C.** Asynchronously cycling p21^{+/+} and p21^{-/-} keratinocytes were suspended in methylcellulose in the presence or absence of 100 μM 5,6-dichloro-1-β-D-ribofuranosyl-benzimidazole (DRB) and harvested at the times indicated. Amounts of cyclin A mRNA and L32 mRNA (loading control) were determined by RNase protection assay.

Submaximal Accumulation of p27^{Kip1} in Suspended p21^{Cip1}-Null Keratinocytes

We have shown that amounts of p27^{Kip1} increase in suspended keratinocytes, as do amounts of p27^{Kip1}-associated cyclin A-cdk2 complexes (18). As shown here, loss of p21^{Cip1} prevents the rapid decline in cyclin A-associated activity in suspended cells. Together, these findings indicate that p27^{Kip1} contributes to cdk2 inactivation in suspended cells but does not compensate for the absence of p21^{Cip1}. As a possible explanation of this phenomenon, we examined p27^{Kip1} expression in p21^{+/+} and p21^{-/-} keratinocytes in methylcellulose.

Amounts of p27^{Kip1} increased substantially when p21^{+/+} keratinocytes were placed in suspension (Fig. 3A); (18). Increases occurred within 6 h of suspension and amounts of p27^{Kip1} remained elevated for at least 24 h. p27^{Kip1} was less abundant in p21^{-/-} keratinocytes than in p21^{+/+} keratinocytes at 0 h (*i.e.*, before detachment) and did not accumulate appreciably in p21^{-/-} keratinocytes suspended for 12 h; submaximal increases were observed at 24 h. Unlike protein, amounts of p27^{Kip1} mRNA were similar in attached and suspended p21^{+/+} and p21^{-/-} keratinocytes (Fig. 3B). These findings show that p21^{Cip1} elicits events that increase the




FIGURE 2. Amounts of cyclin A in suspended p21^{-/-} keratinocytes ectopically expressing p21^{Cip1}. Proliferating p21^{-/-} keratinocytes were infected with adenovirus without insert (Ad) or Ad-p21^{Cip1} (p21) at a multiplicity of infection (MOI) of 250. Cells were suspended in methylcellulose 19 h after infection and harvested 19 h later for determination of p21^{Cip1} and cyclin A abundance by Western blotting. Uninfected p21^{+/+} and p21^{-/-} keratinocytes were also suspended in methylcellulose for 19 h. The p21^{Cip1} antibody used for Western blotting recognizes ectopic (human) p21^{Cip1} but not endogenous (mouse) p21^{Cip1}. A, attached cells; S, suspended cells.




FIGURE 3. Amounts of p27^{Kip1} in suspended wild-type keratinocytes. **A** and **B**. Exponentially growing p21^{+/+} and p21^{-/-} keratinocytes were suspended in methylcellulose and harvested at the times indicated. **A**. Amounts of p27^{Kip1} protein were determined by Western blotting. **B**. Amounts of p27^{Kip1} mRNA and L32 mRNA (loading control) were determined by RNase protection assay.

translation and/or stability of p27^{Kip1} in both adherent and nonadherent keratinocytes. In p21^{-/-} keratinocytes, amounts of p27^{Kip1} are subnormal, and as a result, the potential capacity of p27^{Kip1} to compensate for the absence of p21^{Cip1} is attenuated. Similar to our findings, loss of p21^{Cip1} reduced the accumulation of p27^{Kip1} protein (but not of p27^{Kip1} mRNA) in interleukin-treated astrocytoma cells (28).

Increased Association of CDK Inhibitors With Cyclin A Complexes in Suspended Wild-Type Keratinocytes

In contrast to p27^{Kip1}, p21^{Cip1} did not accumulate in suspended p21^{+/+} keratinocytes (Fig. 4A). In fact, p21^{Cip1} decreased in abundance 6–9 h after suspension and was not detected at 24 h. Although total amounts of p21^{Cip1} were constant for at least 6 h after suspension, amounts of p21^{Cip1} not sequestered by D cyclin-cdk4 complexes (and thus available for interaction with cdk2 complexes) may increase. Consistent with this hypothesis, we have shown that amounts of cyclin D1 decrease after suspension of wild-type keratinocytes (18). To test this hypothesis, we determined the amounts of p21^{Cip1}-associated cyclin A and p21^{Cip1}-free cyclin A in p21^{+/+} keratinocytes at various times after suspension. Cell extracts were immunoprecipitated with antibody to p21^{Cip1}, and p21^{Cip1} immune complexes were Western blotted with antibody to cyclin A (p21^{Cip1}-associated cyclin A). p21^{Cip1}-immunodepleted extracts were immunoprecipitated and Western blotted with antibody to cyclin A (p21^{Cip1}-free cyclin A). Western blots using p21^{Cip1} antibody confirmed that immunodepleted extracts contained little if any p21^{Cip1} (data not shown).

Suspension of p21^{+/+} keratinocytes for 3 or 6 h resulted in an increase in the amounts of p21^{Cip1}-associated cyclin A and a decrease in the amounts of p21^{Cip1}-free cyclin A (Fig. 4B, compare lanes 2 and 3 with lanes 7 and 8). Thus, a greater percentage of cyclin A complexes contain p21^{Cip1} in suspended keratinocytes than in attached keratinocytes. After 6 h, amounts of p21^{Cip1}-associated cyclin A declined (lanes 4 and 5), most

likely because of reductions in p21^{Cip1} abundance (see Fig. 4A). Although suspension reduced amounts of p21^{Cip1}-free cyclin A, it did not eliminate p21^{Cip1}-free cyclin A, at least for times up to 12 h (lane 10). On the other hand, little if any cyclin A remained in 12-h cell extracts immunodepleted of both p21^{Cip1} and p27^{Kip1} (Fig. 4C, top panel, lane 8). Comparison of the relative amounts of CDK inhibitor (CKI)-associated and CKI-free cyclin A at all time points shows that the ratio of associated to free increases progressively after suspension (Fig. 4C, top panel).

The data in Fig. 4 support our hypothesis that suspension-induced decreases in cyclin D1 result in the redistribution of p21^{Cip1} from cyclin D1 complexes to cyclin A complexes and thus in the inhibition of cyclin A-cdk2 activity in wild-type keratinocytes. Redistribution of p27^{Kip1} and/or increases in p27^{Kip1} abundance also contribute to cyclin A-cdk2 inactivation in suspended p21^{+/+} cells. Our studies also explain why cyclin A-cdk2 complexes are active in suspended p21^{-/-} cells: most of these complexes lack p27^{Kip1}, as do most of the cyclin A-cdk2 complexes in attached p21^{-/-} cells. This finding is consistent with our data showing subnormal amounts of p27^{Kip1} in p21^{-/-} keratinocytes (Fig. 3A).




FIGURE 4. Association of cyclin A complexes with p21^{Cip1} and p27^{Kip1} in attached and suspended keratinocytes. **A**. Asynchronously cycling p21^{+/+} keratinocytes were suspended in methylcellulose and harvested at the times indicated. Cell extracts were immunoblotted with antibody to mouse p21^{Cip1}. **B**. Cycling p21^{+/+} keratinocytes were suspended in methylcellulose for the times indicated. Lanes 1–4, cell extracts were immunoprecipitated with antibody to p21^{Cip1}, and p21^{Cip1} immune complexes were Western blotted with antibody to cyclin A. Lanes 5–8, after removal of p21^{Cip1}, cell extracts were immunoprecipitated with antibody to cyclin A, and cyclin A immune complexes were Western blotted with antibody to cyclin A. **C**. As in **B**, with the exception that cell extracts were immunoprecipitated with antibody to both p21^{Cip1} and p27^{Kip1} (p21^{+/+}) or with antibody to p27^{Kip1} (p21^{-/-}).

Reduced Capacity of p21^{-/-} Keratinocytes to Growth Arrest and Differentiate in Suspension

Although suspension typically arrests cells in G₁ (1), suspended primary mouse keratinocytes arrest in G₁, S, and G₂-M (29). To determine whether cell cycle arrest requires p21^{Cip1}, p21^{+/+} and p21^{-/-} keratinocytes attached to tissue culture plates received bromodeoxyuridine (BrdUrd) for 2 h to label S-phase cells. Cells were then suspended in methylcellulose in the absence of BrdUrd and harvested 12 h later, and the position of the BrdUrd-labeled cells in the cell cycle was determined by flow cytometry. The percentages of cells remaining in S phase 12 h after suspension were 52% and 31% for p21^{+/+} and p21^{-/-} populations, respectively (Fig. 5). Moreover, a greater percentage of p21^{-/-} cells were in G₁ at this time than were p21^{+/+} cells. These data suggest that p21^{-/-} cells cycle for longer periods of time when placed in suspension than do p21^{+/+} cells. In this respect, p21^{-/-} keratinocytes mimic MK cells, which are immortalized mouse keratinocytes (30) and which express little if any p21^{Cip1} (data not shown).

To assess the need for p21^{Cip1} for differentiation, we placed p21^{+/+} and p21^{-/-} keratinocytes in suspension, and amounts of two differentiation markers, K1 (Fig. 6A) and loricrin (Fig. 6B), were determined. Both proteins increased in abundance after suspension of p21^{+/+} keratinocytes. Increases in the abundance of loricrin were especially pronounced. Amounts of loricrin did not increase in suspended p21^{-/-} keratinocytes, whereas amounts of K1 increased to a lesser extent and in a delayed manner. At 48 h, K1 was still much less abundant in p21^{-/-} keratinocytes than in p21^{+/+} keratinocytes (data not shown).

These findings show that efficient differentiation of primary keratinocytes requires p21^{Cip1}. To confirm that the differentiation-defective phenotype of p21^{-/-} keratinocytes reflects the absence of p21^{Cip1}, p21^{-/-} keratinocytes were infected with Ad-p21^{Cip1} and suspended in methylcellulose. Nineteen hours after suspension, Ad-p21^{Cip1}-infected p21^{-/-} cells (Fig. 6C, lane 4) expressed more K1 than did uninfected p21^{-/-} cells (lane 2) and p21^{-/-} cells infected with adenovirus alone (lane 3).




FIGURE 5. Proliferative capacity of suspended wild-type and p21^{Cip1}-null keratinocytes. Cycling p21^{+/+} and p21^{-/-} cells received 4 μM BrdUrd for 2 h. Cell pellets were processed as described in "Materials and Methods," and cell cycle distribution was determined by flow cytometry. Percentages of BrdUrd-labeled cells in G₁, S, and G₂ are shown.




FIGURE 6. K1 and loricrin induction in p21^{+/+} and p21^{-/-} keratinocytes suspended in methylcellulose. **A** and **B.** p21^{+/+} and p21^{-/-} keratinocytes were suspended in methylcellulose for the indicated times, and cell extracts were immunoblotted with antibody to K1 (**A**) or loricrin (**B**). **C.** p21^{-/-} keratinocytes were infected with control adenovirus (*Ad*) or Ad-p21^{Cip1} (*p21*) at an MOI of 250. Cells were suspended in methylcellulose 21 h after infection and harvested 19 h later for determination of K1 abundance by Western blotting. Lane 1, uninfected cells attached to plates.

E2F Complexes in Wild-Type and p21^{Cip1}-Null Keratinocytes in Suspension

Cyclin A is a member of a DNA binding complex that contains cdk2, the transcription factor E2F, the E2F binding partner DP1, and p107, an Rb family member (31–33). Previous studies have shown that p21^{Cip1} disrupts this complex both *in vivo* and *in vitro* (21, 34, 35). Thus, as another potential consequence of p21^{Cip1} loss, we monitored E2F DNA binding activity in p21^{+/+} and p21^{-/-} keratinocytes before and after suspension. As demonstrated by electrophoretic mobility shift assays, attached keratinocytes (wild-type and p21^{Cip1}-null) expressed an activity that interacted with an E2F binding site; this DNA/protein complex is designated complex I (Fig. 7A, lanes 1 and 5). Between 3 and 6 h after suspension of p21^{+/+} keratinocytes, complex I disappeared and a more rapidly migrating complex (complex II) appeared (Fig. 7A, lanes 3 and 4). Complex II was not seen in suspended p21^{-/-} keratinocytes, although amounts of complex I decreased somewhat (lanes 6–8). Infection of p21^{-/-} keratinocytes with Ad-p21^{Cip1} resulted in the conversion of complex I to II (Fig. 7B). These findings show that suspension modulates the composition of E2F DNA binding complexes in a p21^{Cip1}-dependent manner.

Antibody to cyclin A blocked the formation of complex I (or its entry into the gel; Fig. 7C, compare lanes 1 and 4, and lanes 7–9 with lanes 10–12) but had no effect on complex




FIGURE 7. Composition of E2F DNA binding complexes in suspended p21^{+/+} and p21^{-/-} keratinocytes. **A–D.** Electrophoretic mobility shift assays were performed on cell extracts using a fragment of the dihydrofolate reductase promoter containing an E2F binding site. Antibodies to cyclin A (**C**), E2F1, E2F2, E2F3, E2F4, E2F5, p130, p107, or Rb (**D**) were added to binding reactions 30 min before addition of the radiolabeled DNA probe. **A.** p21^{+/+} and p21^{-/-} keratinocytes were suspended for the indicated times. **B.** p21^{Cip1}-null keratinocytes were infected with control adenovirus (*Ad*) or Ad-p21^{Cip1} (*p21*) at an MOI of 250. Cells were suspended 19 h after infection and harvested 9 h later. Uninfected wild-type keratinocytes were also suspended for 9 h. **C.** p21^{+/+} and p21^{-/-} keratinocytes were suspended for the indicated times. **D.** p21^{-/-} keratinocytes (*top panel*) and p21^{+/+} keratinocytes (*bottom panel*) were suspended for 6 h. The *top panel* examines the composition of complex I, and the *bottom panel* examines the composition of complex II.

II (compare *lanes 2* and *3* with *lanes 5* and *6*). Both complexes were supershifted with antibodies to p107 and E2F4 but were unaffected by antibodies to Rb, the Rb-related protein p130, or other E2F family members (Fig. 7D). Thus, complex I contains E2F4, p107, and cyclin A, whereas complex II contains E2F4 and p107 but not cyclin A.

Discussion

Our data show that suspension reduces the abundance and associated activity of cyclin A in primary keratinocytes much more rapidly in the presence than in the absence of p21^{Cip1}. Suspension-induced decreases in cyclin A abundance apparently result (at least in part) from the transcriptional repression of the *cyclin A* gene. In wild-type keratinocytes, suspension reduced the abundance of the cyclin A transcript without affecting its half-life. This transcript also decayed with similar kinetics in p21^{+/+} and p21^{-/-} keratinocytes. Others have shown that suspension represses cyclin A expression in rodent fibroblasts by a process involving cdk2 inactivation and the consequent interaction of Rb with the cyclin A promoter (10, 14, 21, 22). In our studies, loss of cyclin A activity preceded loss of cyclin A protein. Thus, the primary target of p21^{Cip1} in suspended keratinocytes is cyclin A-associated activity rather than cyclin A expression.

We show that p21^{Cip1} inhibits cyclin A-associated activity in suspended keratinocytes by two mechanisms: it interacts with cyclin A-cdk2 complexes, and it promotes the accumulation of p27^{Kip1}. Suspension of p21^{+/+} keratinocytes for 3 or 6 h increased the percentage of cyclin A complexes containing p21^{Cip1}. Such increases presumably result from the redistribution of p21^{Cip1} from cyclin D1-cdk4 complexes to cyclin A-cdk2 complexes. First, as presented here, amounts of p21^{Cip1} remained constant for 6–9 h after suspension of wild-type keratinocytes and then decreased. Second, as described previously, amounts of cyclin D1 declined within 3 h of suspension of wild-type keratinocytes and were essentially undetectable at 6 h (18). p21^{Cip1} may interact with both cyclin A-cdk2 and cyclin E-cdk2 complexes in suspended keratinocytes; we have shown that suspension reduces cyclin E-associated activity and cyclin A-associated activity with similar kinetics (18). Others have shown that p21^{Cip1} associates with cyclin E-cdk2 complexes in suspended NIH 3T3 cells and that this association correlates with decreases in cyclin D1 abundance rather than increases in p21^{Cip1} abundance (9).

Amounts of p27^{Kip1} increased within 6 h of suspension of wild-type keratinocytes and were substantially elevated at 9–24 h. p21^{+/+} keratinocytes expressed much more p27^{Kip1} than did p21^{-/-} keratinocytes whether attached or suspended.

Consistent with these results, only a small percentage of cyclin A complexes contained p27^{Kip1} in suspended p21^{-/-} keratinocytes. Immunodepletion of p21^{Cip1} from extracts of p21^{+/+} cells suspended for 6 or 12 h partially removed cyclin A complexes, whereas immunodepletion of both p21^{Cip1} and p27^{Kip1} removed almost all cyclin A complexes. Thus, both CKIs contribute to the formation of inactive cyclin A-cdk2 complexes in suspended wild-type keratinocytes. In this regard, p21^{Cip1} may be important at early times after suspension (*i.e.*, at times before p27^{Kip1} increases in abundance), whereas p27^{Kip1} is important at later times (*i.e.*, at times when p27^{Kip1} is abundant and p21^{Cip1} is not appreciably expressed). The data of Di Cunto *et al.* (36) show that enforced expression of p21^{Cip1} blocks the terminal stages of keratinocyte differentiation. Thus, the “p21^{Cip1} to p27^{Kip1} switch” may represent a means of keeping cdk2 inactive while allowing the completion of differentiation.

How p21^{Cip1} modulates p27^{Kip1} abundance in keratinocytes is not known. p21^{+/+} and p21^{-/-} keratinocytes express comparable amounts of p27^{Kip1} mRNA whether attached or suspended. cdk2 phosphorylates p27^{Kip1}, which targets p27^{Kip1} for destruction in the proteasome (37–39). Thus, the greater amounts of p27^{Kip1} in p21^{+/+} than in p21^{-/-} keratinocytes may be the indirect result of reduced amounts of cdk2 activity and consequent p27^{Kip1} stabilization. On the other hand, Carrano and Pagano (16) found that suspension of cycling fibroblasts accelerated p27^{Kip1} degradation in the absence of cdk2 activity; whether cdk2-independent p27^{Kip1} destruction required p21^{Cip1} was not determined.

We have shown that wild-type keratinocytes withdraw from the cell cycle regardless of whether they are in G₁, S, or G₂-M at the time of suspension (29). Results of experiments using BrdUrd to tag S phase cells show that more p21^{-/-} cells exit S phase when placed in suspension than do p21^{+/+} cells. Thus, loss of p21^{Cip1} appears to increase the capacity of keratinocytes to proliferate in suspension. Using a similar method, others (40) showed that ectopic expression of p21^{Cip1} prevented SAOS-2 cells from exiting S phase. p21^{Cip1} may arrest suspension-cultured keratinocytes in S phase arrest by inhibiting the activity of cdk2, as was shown in the SAOS-2 study (40), and/or by directly interfering with DNA replication (41).

p21^{Cip1}-null mice develop normally (42), thus indicating that p21^{Cip1} is not required for keratinocyte differentiation during development. However, our data suggest that loss of p21^{Cip1} impedes differentiation to some extent in suspension-cultured keratinocytes. We show that suspended p21^{-/-} keratinocytes express much less K1, a marker of differentiation, than do suspended p21^{+/+} keratinocytes or p21^{-/-} keratinocytes ectopically expressing p21^{Cip1}. A second differentiation marker, loricrin, increased in suspended p21^{+/+} keratinocytes but not in p21^{-/-} keratinocytes. The differentiation-defective phenotype of p21^{-/-} keratinocytes may reflect their impaired capacity to growth arrest in suspension and/or to accumulate p27^{Kip1}. We have shown that keratinocytes treated with p27^{Kip1} antisense oligonucleotides or derived from p27^{Kip1}-null mice do not differentiate when placed in suspension (18, 43). Similar to our results, Missero *et al.* (44) found that loss of p21^{Cip1} reduced the expression of a set of differentiation markers in calcium-treated mouse keratinocytes.

E2F-p107 complexes associate with cyclin E-cdk2 and cyclin A-cdk2 complexes in G₁ and S, respectively (31–33, 45). Our data show that suspension of wild-type keratinocytes results in the removal of cyclin A from E2F-p107 DNA binding complexes, whereas suspension of p21^{Cip1}-null keratinocytes does not. Similarly, others have shown that cyclin E and cdk2 associate with E2F-p107 complexes in attached but not suspended NIH 3T3 cells (10, 19) and that p21^{Cip1} releases cyclin-cdk2 complexes from E2F-p107 complexes *in vitro* (34, 35). The physiological relevance of cyclin-cdk2 interaction with E2F-p107 complexes is not known at present (46). However, it is clear that among its other effects, p21^{Cip1} modulates the composition of E2F-p107 complexes in primary keratinocytes.

Although the p21^{Cip1} gene is rarely mutated in human tumors (47), p21^{Cip1} may be expressed at subnormal amounts or dysfunctional in some human tumors (48, 49). Importantly, inefficient expression of p21^{Cip1} is indicative of short survival times for patients with acute lymphoblastic leukemia or colorectal cancer (48, 50). We have shown that irradiated p21^{Cip1}-null mice develop significantly more metastatic tumors than do irradiated wild-type mice (51). A key step in the metastatic process is the loss of adherent-dependent growth control (52). Thus, loss of p21^{Cip1} may contribute to tumor progression by increasing the capacity of cells to grow in suspension.

Materials and Methods

Cell Culture

p21^{Cip1}-null mice (from Dr. Tyler Jacks; 53) and p21^{Cip1}-wild type mice (The Jackson Laboratory, Bar Harbor, ME), both from the same mixed C57BL/6 and 129 genetic background, were used in our study. Keratinocytes were isolated from the epidermis of newborn mice as described previously (25, 54) and cultured in keratinocyte basal medium (KBM; Clonetics Corp., San Diego, CA) supplemented with 0.05 mM calcium, 0.5 μg/ml insulin, 0.05 μg/ml hydrocortisone, 1.5 ng/ml epidermal growth factor, 7.5 μg/ml bovine pituitary extract, and 2% dialyzed FCS. Confluent 3-day-old keratinocytes were refed with KBM containing the above ingredients 12–15 h before suspension. Cells were suspended in 50-ml conical tubes in KBM containing 1.45% methylcellulose (Sigma, St. Louis, MO) and all of the above ingredients except epidermal growth factor.

Preparation of Cell Extracts and Western Blotting

Adherent keratinocytes were trypsinized from the plates, collected by centrifugation, and washed with KBM containing 5% serum to inactivate the trypsin. Keratinocytes suspended in methylcellulose were diluted 1/10 with PBS and collected by centrifugation. Cells were rinsed with PBS, resuspended in immunoprecipitation buffer [50 mM HEPES (pH 7.2), 150 mM NaCl, 0.1% Tween 20, 1 mM EDTA, 1 mM EGTA, 0.1 mM orthovanadate, 0.5 mM NaF, 0.1 mM phenylmethylsulfonyl fluoride, 2.5 μg/ml leupeptin, and 1 mM DTT], and sonicated for 5 s. Insoluble material was removed by centrifugation. Cell extracts (25–50 μg protein) were resolved by SDS-PAGE and transferred to nitrocellulose membranes. Membranes were

blocked in PBS containing 0.1% Tween 20 and 5% instant milk and incubated with primary antibody in PBS containing 0.1% Tween 20 for 1 h at room temperature. Proteins recognized by the primary antibody were detected by enhanced chemiluminescence using a horseradish peroxidase-coupled secondary antibody as specified by the manufacturer (Amersham Biosciences, Piscataway, NJ).

In Vitro Kinase Assays

Cell extracts (20–40 μg of protein) were incubated with antibody to cyclin A for 2–10 h at 4°C and subsequently with protein A-agarose beads for 30 min at 4°C. Immune complexes were collected by centrifugation and washed twice with immunoprecipitation buffer and once with kinase buffer [50 mM HEPES (pH 7.5) and 10 mM MgCl_2]. Washed complexes were incubated for 5 min at 30°C in kinase buffer containing 0.5 μCi [γ - ^{32}P]ATP, 1 μM ATP, and 100 $\mu\text{g}/\text{ml}$ histone H1. Reaction mixtures were applied to SDS-polyacrylamide gels, and phosphoproteins were visualized by autoradiography.

Detection of Cells Labeled With BrdUrd

Cells received 4 μM BrdUrd for 2 h. Rinsed cell pellets were fixed in 60% cold ethanol and incubated in 0.04% pepsin (5 ml in 0.1 N HCl) for 40 min at 37°C. Cells were centrifuged, resuspended in 3 ml of 2 N HCl, and incubated for 20 min at 37°C. Cells were pelleted after addition of 6 ml of 0.1 N sodium borate to the samples, washed with PBS containing 0.1% Tween 20, and incubated in the dark for 1 h at room temperature in 0.2 ml of PBS containing 0.1% Tween 20 and 0.4 μl of BrdUrd-FITC antibody (Boehringer-Mannheim, Indianapolis, IN). Cells were washed twice with PBS containing 0.1% Tween 20 and incubated for 25 min at room temperature in PBS containing 0.1% Tween 20, 0.05% BSA, 20 $\mu\text{g}/\text{ml}$ propidium iodide, and 50 $\mu\text{g}/\text{ml}$ RNase. Cell cycle distributions were determined using a Becton Dickinson FACScan.

Electrophoretic Mobility Shift Assays

E2F DNA-binding assays were performed essentially as described by Ikeda *et al.* (55). Cell pellets were incubated on ice for 30 min in a buffer containing 50 mM HEPES (pH 7.9), 250 mM KCl, 0.1 mM EDTA, 0.1 mM EGTA, 0.1% NP40, 0.4 mM NaF, 0.4 mM vanadate, 10% glycerol, 1 mM DTT, 0.5 mM phenylmethylsulfonyl fluoride, and 2.5 $\mu\text{g}/\text{ml}$ leupeptin. After removal of insoluble material, extracts (20 μg of protein) were incubated for 30 min at room temperature in binding buffer [20 mM HEPES (pH 7.9), 40 mM KCl, 6 mM MgCl_2 , 1 mM EGTA, 1 mM DTT, 0.1% NP40, 30 mg/ml BSA, and 500 ng/ml sonicated salmon sperm DNA] containing a ^{32}P -labeled DNA fragment (0.1 ng) corresponding to residues –103 to –23 of the dihydrofolate reductase promoter in a final volume of 50 μl . DNA-protein complexes were separated on 5% non-denaturing polyacrylamide gels (75:1 acrylamide/bisacrylamide) in Tris-borate with 5% glycerol and visualized by autoradiography. For supershifts, extracts were incubated with antibodies for 30 min before addition of the radiolabeled DNA probe.

Adenovirus Infection

Stocks of adenovirus encoding green fluorescent protein with and without p21^{Cip1} were produced, purified, and titered as described previously (56, 57). Adenoviruses were provided by Dr. Joseph Nevins at Duke University.

Messenger RNA Analysis

Total mRNA was isolated using Trizol Reagent (Invitrogen, Carlsbad, CA). For RNase protection assays, mRNA (20 μg) was hybridized overnight at 56°C with ^{32}P -labeled probes (10⁵ cpm) from a custom-made probe set (PharMingen, San Diego, CA). Samples were digested with RNase T1 and RNase A for 45 min at 30°C and proteinase K for 15 min at 37°C. Samples were extracted with phenol/chloroform, collected by sodium acetate/ethanol precipitation, denatured at 90°C for 3 min, and electrophoresed on a 5% polyacrylamide gel. Gels were dried and exposed to X-ray film.

Acknowledgments

The authors thank Dr. Tyler Jacks for the p21^{Cip1}-null mice, Dr. Joseph Nevins for the adenoviruses, and Dr. Nancy Olashaw for manuscript preparation. We also acknowledge the helpful service of the Molecular Biology, Molecular Imaging, and Flow Cytometry Core Laboratories at the Moffitt Cancer Center.

References

1. Assoian RK. Control of the G1 phase cyclin-dependent kinases by mitogenic growth factors and the extracellular matrix. *Cytokine Growth Factor Rev*, 1997;8:165–70.
2. Olashaw N, Pledger WJ. Paradigms of growth control: relation to Cdk activation. *Science STKE*, 2002;2002:RE7.
3. Weinberg RA. The retinoblastoma protein and cell cycle control. *Cell*, 1995; 81:323–30.
4. DeGregori J, Kowalik T, Nevins JR. Cellular targets for activation by the E2F1 transcription factor include DNA synthesis and G1/S regulatory genes. *Mol Cell Biol*, 1995;15:4215–24.
5. Lundberg AS, Weinberg RA. Functional inactivation of the retinoblastoma protein requires sequential modification by at least two distinct cyclin-cdk complexes. *Mol Cell Biol*, 1998;18:753–61.
6. Sherr CJ, Roberts JM. Inhibitors of mammalian G1 cyclin-dependent kinases. *Genes Dev*, 1995;9:1149–63.
7. Bagui TK, Jackson RJ, Agrawal D, Pledger WJ. Analysis of cyclin D3-cdk4 complexes in fibroblasts expressing and lacking p27^{Kip1} and p21^{Cip1}. *Mol Cell Biol*, 2000;20:8748–87.
8. Guadagno TM, Ohtsubo M, Roberts JM, Assoian RK. A link between cyclin A expression and adhesion-dependent cell cycle progression. *Science*, 1993;262:1572–5.
9. Zhu X, Ohtsubo M, Bohmer RM, Roberts JM, Assoian RK. Adhesion-dependent cell cycle progression linked to the expression of cyclin D1, activation of cyclin E-cdk2, and phosphorylation of the retinoblastoma protein. *J Cell Biol*, 1996;133:391–403.
10. Schulze A, Zerfass-Thome K, Berges J, Middendorp S, Jansen-Durr P, Henglein B. Anchorage-dependent transcription of the cyclin A gene. *Mol Cell Biol*, 1996;16:4632–8.
11. Kang JS, Krauss RS. Ras induces anchorage-independent growth by subverting multiple adhesion-regulated cell cycle events. *Mol Cell Biol*, 1996;16:3370–80.
12. Bottazzi ME, Zhu X, Bohmer RM, Assoian RK. Regulation of p21^{Cip1} expression by growth factors and the extracellular matrix reveals a role for transient ERK activity in G1 phase. *J Cell Biol*, 1999;146:1255–64.
13. Barrett JF, Lewis BC, Hoang AT, Alvarez RJ Jr, Dang CV. Cyclin A links c-Myc to adhesion-independent cell proliferation. *J Biol Chem*, 1995;270: 15923–5.
14. Philips A, Huet X, Plet A, Rech J, Vie A, Blanchard JM. Anchorage-dependent expression of cyclin A in primary cells requires a negative DNA regulatory element and a functional Rb. *Oncogene*, 1999;18:1819–25.
15. Carstens CP, Kramer A, Fahl WE. Adhesion-dependent control of cyclin

- E/cdk2 activity and cell cycle progression in normal cells but not in Ha-ras transformed NRK cells. *Exp Cell Res*, 1996;229:86–92.
16. Carrano AC, Pagano M. Role of the F-box protein Skp2 in adhesion-dependent cell cycle progression. *J Cell Biol*, 2001;153:1381–90.
 17. Radeva G., Petrocelli T, Behrend E, Leung-Hagetejin C, Filmus J, Slingerland J, et al. Overexpression of the integrin-linked kinase promotes anchorage-independent cell cycle progression. *J Biol Chem*, 1997;272:13937–44.
 18. Hauser PJ, Agrawal D, Flanagan M, Pledger WJ. The role of p27^{Kip1} in the *in vitro* differentiation of murine keratinocytes. *Cell Growth & Differ*, 1997;8:203–11.
 19. Schulze A, Mannhardt B, Zerfass-Thome K, Zwerschke W, Jansen-Durr P. Anchorage-independent transcription of the cyclin A gene induced by the E7 oncoprotein of human papillomavirus type 16. *J Virol*, 1998;72:2323–34.
 20. Resnitzky D. Ectopic expression of cyclin D1 but not cyclin E induces anchorage-independent cell cycle progression. *Mol Cell Biol*, 1997;17:5640–47.
 21. Zerfass-Thome K, Schulze A, Zwerschke W, Vogt B, Helin K, Bartek J, et al. p27^{Kip1} blocks cyclin E-dependent transactivation of cyclin A gene expression. *Mol Cell Biol*, 1997;17:407–15.
 22. Kramer A, Carstens CP, Fahl WE. A novel CCAAT-binding protein necessary for adhesion-dependent cyclin A transcription at the G1/S boundary is sequestered by a retinoblastoma-like protein in G0. *J Biol Chem*, 1996;271:6579–82.
 23. Fang F, Orend G, Watanabe N, Hunter T, Ruoslahti E. Dependence of cyclin E-CDK2 kinase activity on cell anchorage. *Science*, 1996;271:499–502.
 24. Kawada M, Yamagoe S, Murakami Y, Suzuki K, Mizuno S, Uehara Y. Induction of p27^{Kip1} degradation and anchorage independence by Ras through the MAP kinase signaling pathway. *Oncogene*, 1997;15:629–37.
 25. Drozdoff V, Pledger WJ. Commitment to differentiation and expression of early differentiation markers in murine keratinocytes *in vitro* are regulated independently of extracellular calcium concentrations. *J Cell Biol*, 1993;123:909–19.
 26. Pagano M, Pepperkok R, Verde F, Ansorge W, Draetta G. Cyclin A is required at two points in the human cell cycle. *EMBO J*, 1992;11:961–71.
 27. Harper JW, Elledge SJ, Keyomarsi K, Dynlacht B, Tsai LH, Zhang P, et al. Inhibition of cyclin-dependent kinases by p21. *Mol Biol Cell*, 1995;6:387–400.
 28. Liu J, Estes ML, Drazba JA, Liu H, Prayson R, Kondo S, et al. Anti-sense oligonucleotide of p21^{waf1/cip1} prevents interleukin 4-mediated elevation of p27^{Kip1} in low grade astrocytoma cells. *Oncogene*, 2000;19:661–9.
 29. Hauser PJ, Agrawal D, Pledger WJ. Primary keratinocytes have an adhesion dependent S phase checkpoint that is absent in immortalized cell lines. *Oncogene*, 1998;17:3083–92.
 30. Weissman BE, Aaronson SA. BALB and Kirsten murine sarcoma viruses alter growth and differentiation of EGF-dependent balb/c mouse epidermal keratinocyte lines. *Cell*, 1983;32:599–606.
 31. Cao L, Faha B, Dembski M, Tsai LH, Harlow E, Dyson N. Independent binding of the retinoblastoma protein and p107 to the transcription factor E2F. *Nature*, 1992;355:176–9.
 32. Devoto SH, Mudryj M, Pines J, Hunter T, Nevins JR. A cyclin A-protein kinase complex possesses sequence-specific DNA binding activity: p33^{cdk2} is a component of the E2F-cyclin A complex. *Cell*, 1992;68:167–76.
 33. Shirodkar S, Ewen M, DeCaprio JA, Morgan J, Livingston DM, Chittenden T. The transcription factor E2F interacts with the retinoblastoma product and a p107-cyclin A complex in a cell cycle-regulated manner. *Cell*, 1992;68:157–66.
 34. Zhu L, Harlow E, Dynlacht BD. p107 uses a p21^{Cip1}-related domain to bind cyclin/cdk2 and regulate interactions with E2F. *Genes Dev*, 1995;9:1740–52.
 35. Afshari CA, Nichols MA, Xiong Y, Mudryj M. A role for a p21-E2F interaction during senescence arrest of normal human fibroblasts. *Cell Growth & Differ*, 1996;7:979–88.
 36. Di Cunto F, Topley G, Calautti E, Hsiao J, Ong L, Seth PK, et al. Inhibitory function of p21^{Cip1/WAF1} in differentiation of primary mouse keratinocytes independent of cell cycle control. *Science*, 1998;280:1069–72.
 37. Sheaff RJ, Groudine M, Gordon M, Roberts JM, Clurman BE. Cyclin E-CDK2 is a regulator of p27^{Kip1}. *Genes Dev*, 1997;11:1464–78.
 38. Vlach J, Hennecke S, Amati B. Phosphorylation-dependent degradation of the cyclin-dependent kinase inhibitor p27. *EMBO J*, 1997;16:5334–44.
 39. Morisaki H, Fujimoto A, Ando A, Nagata Y, Ikeda K, Nakanishi M. Cell cycle-dependent phosphorylation of p27 cyclin-dependent kinase (Cdk) inhibitor by cyclin E/Cdk2. *Biochem Biophys Res Commun*, 1997;240:386–90.
 40. Ogryzko VV, Wong P, Howard BH. WAF1 retards S-phase progression primarily by inhibition of cyclin-dependent kinases. *Mol Cell Biol*, 1997;17:4877–82.
 41. Waga S, Hannon GJ, Beach D, Stillman B. The p21 inhibitor of cyclin-dependent kinases controls DNA replication by interaction with PCNA. *Nature*, 1994;369:574–8.
 42. Deng C, Zhang P, Harper JW, Elledge SJ, Leder P. Mice lacking p21^{Cip1/WAF1} undergo normal development, but are defective in G1 checkpoint control. *Cell*, 1995;82:675–84.
 43. Hauser PJ, Agrawal D, Hackney J, Pledger WJ. STAT3 activation accompanies keratinocyte differentiation. *Cell Growth & Differ*, 1998;9:847–55.
 44. Missero C, Di Cunto F, Kiyokawa H, Koff A, Dotto GP. The absence of p21^{Cip1/WAF1} alters keratinocyte growth and differentiation and promotes rasi-tumor progression. *Genes Dev*, 1996;10:3065–75.
 45. Lees E, Faha B, Dulic V, Reed SI, Harlow E. Cyclin E/cdk2 and cyclin A/cdk2 kinases associate with p107 and E2F in a temporally distinct manner. *Genes Dev*, 1992;6:1874–85.
 46. Smith EJ, Nevins JR. The Rb-related p107 protein can suppress E2F function independently of binding to cyclin A/cdk2. *Mol Cell Biol*, 1995;15:338–44.
 47. Shiohara M, Koike K, Komiyama A, Koeffler HP. p21^{waf1} mutations and human malignancies. *Leuk Lymphoma*, 1997;26:35–41.
 48. Roman-Gomez J, Castillejo JA, Jimenez A, Gonzalez MG, Moreno F, Rodriguez MDCC, et al. 5' CpG island hypermethylation is associated with transcriptional silencing of the p21^{Cip1/WAF1/Sd11} gene and confers poor prognosis in acute lymphoblastic leukemia. *Blood*, 2002;99:2291–6.
 49. Zhou BP, Liao Y, Xia W, Spohn B, Lee MH, Hung MC. Cytoplasmic localization of p21^{Cip1/WAF1} by Akt-induced phosphorylation in HER-2/*neu*-overexpressing cells. *Nat Cell Biol*, 2001;3:245–52.
 50. Zirbes TK, Baldus SE, Moenig SP, Nolden S, Kunze D, Shafizadeh ST, et al. Prognostic impact of p21^{waf1/cip1} in colorectal cancer. *Int J Cancer*, 2000;89:14–18.
 51. Jackson RJ, Engelman RW, Coppola D, Cantor AB, Wharton W, Pledger WJ. p21^{Cip1} nullizygosity increases tumor metastasis in irradiated mice. *Cancer Res*, 2003;63:3021–5.
 52. Bashyam MD. Understanding cancer metastasis: an urgent need for using differential gene expression analysis. *Cancer*, 2002;94:1821–9.
 53. Brugarolas J, Chandrasekaran C, Gordon JI, Beach D, Jacks T, Hannon GJ. Radiation-induced cell cycle arrest compromised by p21 deficiency. *Nature*, 1995;377:552–7.
 54. Yuspa SH, Harris CC. Altered differentiation of mouse epidermal cells treated with retinyl acetate *in vitro*. *Exp Cell Res*, 1974;86:86–105.
 55. Ikeda MA, Jakoi L, Nevins JR. A unique role for the Rb protein in controlling E2F accumulation during cell growth and differentiation. *Proc Natl Acad Sci USA*, 1996;93:3215–20.
 56. Nevins JR, DeGregori J, Jakoi L, Leone G. Functional analysis of E2F transcription factor. *Methods Enzymol*, 1997;283:205–19.
 57. Song L, Turkson J, Karras JG, Jove R, Haura EB. Activation of Stat3 by receptor tyrosine kinases and cytokines regulates survival in human non-small cell carcinoma cells. *Oncogene*, 2003;22:4150–65.

Molecular Cancer Research

Efficient Down-Regulation of Cyclin A-Associated Activity and Expression in Suspended Primary Keratinocytes Requires p21^{Cip1} 1 This work was supported by the Cortner-Couch Endowed Chair for Cancer Research (W. J. P.) and NIH Grants CA73780 (W. J. P.) and CA90489 (W. D. C.)

Paul Hauser, Le Ma, Deepak Agrawal, et al.

Mol Cancer Res 2004;2:96-104.

Updated version Access the most recent version of this article at:
<http://mcr.aacrjournals.org/content/2/2/96>

Cited articles This article cites 55 articles, 33 of which you can access for free at:
<http://mcr.aacrjournals.org/content/2/2/96.full#ref-list-1>

Citing articles This article has been cited by 4 HighWire-hosted articles. Access the articles at:
<http://mcr.aacrjournals.org/content/2/2/96.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://mcr.aacrjournals.org/content/2/2/96>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.